

המחלקה להנדסת חשמל ואלקטרוניקה

תאריך הבחינה : 23.02.17

שעות הבחינה : 9:00-12:00

## מבוא לאותות אקריאים

מועד ב'

ד"ר דימה בחובסקי

תשע"ז סטטוס א'

חומר עזר - עד 4 דפי נוסחאות אישיים (משני צדדים), מחשבון

הוראות מיוחדות :

- השאלון כולל 3 שאלות ללא בחירה, סך הכל של 110 נקודות.
- יש לציין באופן מלא וברור את שלבי הפתרון. תשובה ללא הסבר לא תתקבלנה.
- אם לא מצויין אחרת, הסעיפים הם בעלי ניקוד זהה.

השאלון כולל 10 דפים (כולל דף זה)

בצלחה !

## 1 תהליכיים אקראיים – סטציואונריות (40 נק')

נתונים תהליכיים אקראיים בלתי תלויים, WSS ו-  $X(t)$ ,  $Y(t)$  כך ש-  
 $E[X(t)] = E[Y(t)] = 1$ ,  $Var[X(t)] = \sigma_X^2$ ,  $Var[Y(t)] = \sigma_Y^2$   
 $.V(t) = X(t) + aY(t)$ ,  $W(t) = X(t)Y(t)$  :

1. האם  $V(t)$  WSS הוא ?

2. האם  $W(t)$  WSS הוא ?

3. חשב  $R_{VW}(t_1, t_2)$  או  $R_{VW}(t, t + \tau)$  ? האם מדובר בתהליכיים סטציואונריים במשמעות ?

4. חשב ערך של  $a$  עבורו  $V(t), W(t)$ , בהינתן  $R_x(\tau) = R_y(\tau)$  אורטוגונליים,

## 2 תהליכיים אקראיים – תכונות (40 נק')

נתנו אותן אקראי מהצורה

$$X(t) = A_c \left[ 1 + m \sin(2\pi f_m t + \phi) \right] \cos(2\pi f_c t + \theta)$$

כאשר קבועים  $m, A_c$ ,  $\phi, \theta \sim U[0, 2\pi]$  הם משתנים אקראיים בלתי תלויים.

מצא :

1.  $E[X(t)]$

2.  $Var[X(t)]$

3.  $R_x(t_1, t_2)$  או  $R_x(t, t + \tau)$  ? האם מדובר בתהליך סטציואונרי ?

4. אותן עובר דרך מערכת

$$H(f) = -j \operatorname{sgn}(f) = \begin{cases} -j & f > 0 \\ j & f < 0 \end{cases}$$

חשב הספק  $P_y$  בМОץם המערכת.

### 3 תהליכי Poisson (30 נק')

חלקיים פוגעים בגלאי ממוצע אחד ל-2 שניות. חשב:

1. סיכוי ל-2 חלקיים בדיק בתוך שנייה מסוימת.

2. סיכוי ללא יותר מחלקיק 1 בתוך שנייה מסוימת.

מעוניינים למדוד קצב הפגיעה של הגלאי ממוצע ע"י הגלאי.

נניח תוגבה להלם של הגלאי מהצורה  $a = 10 \left[ \frac{1}{sec} \right]$ , כאשר  $h(t) = e^{-at}u(t)[V]$ , כאשר  $t_i$ ,  $X(t) = \sum_{i=1}^{\infty} h(t-t_i)$ , זה זמן פגיעה החלקיים. תוכלת כתועא מפיגעות החלקיים נוצר אותה מהצורה  $E[X(t)] = 0.1[V]$  אחרי 10 דק' מדידה.

3. מהו קצב ממוצע של פגיעה החלקיים בגלאי?

# Random Processes – Formulas

## 1 Random Variables

### 1.1 Distributions

	Notation	PDF/PMF	CDF	$E[X]$	$\text{Var}[X]$
Uniform	$U[a,b]$	$\frac{1}{b-a}, a \leq x \leq b$	$\begin{cases} 0 & x < a \\ \frac{x-a}{b-a} & a \leq x \leq b \\ 1 & b < x \end{cases}$	$\frac{a+b}{2}$	$\frac{(b-a)^2}{12}$
Normal	$N(\mu, \sigma^2)$	$\frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$	$\Phi(x)$	$\mu$	$\sigma^2$
Exponential	$Exp(\lambda)$	$\lambda e^{-\lambda x}, x \geq 0$	$1 - e^{-\lambda x}$	$1/\lambda$	$1/\lambda^2$
Poisson	$\mathcal{P}(\lambda)$	$p(X = k) = e^{-\lambda} \frac{\lambda^k}{k!}$	$e^{-\lambda} \sum_{i=0}^k \frac{\lambda^i}{i!}$	$\lambda$	$\lambda$
Erlang	$Erlang(k, \lambda t)$	$\lambda \frac{(\lambda t)^{k-1}}{(k-1)!} e^{-\lambda t}$	$1 - \sum_{n=0}^{k-1} \frac{(\lambda t)^n}{n!} e^{-\lambda t}$	$k/\lambda t$	$k/(\lambda t)^2$

Special properties:

- Given sum of two independent distributions  $X \sim \mathcal{P}(\lambda_1)$  and  $Y \sim \mathcal{P}(\lambda_2)$ , the resulting distribution is given by  $X + Y \sim \mathcal{P}(\lambda_1 + \lambda_2)$ .
- If  $X_i \sim Exp(\lambda)$  then  $\sum_{i=1}^k X_i \sim Erlang(k, \lambda)$ .

### 1.2 Properties

Definitions:

$F_X(x) = p(X \leq x)$	(1a)	Expectation
$f_X(x) = \frac{\partial F_X(x)}{\partial x}$	(1b)	$E[X] = \begin{cases} \int_{-\infty}^{\infty} x f_X(x) dx \\ \sum_i x_i p(X = x_i) \end{cases} \quad (2a)$
$F_X(x) = \int_{-\infty}^x f_X(p) dp$	(1c)	$E[g(X)] = \begin{cases} \int_{-\infty}^{\infty} g(x) f_X(x) dx \\ \sum_i g(x_i) p(X = x_i) \end{cases} \quad (2b)$
		$E[aX] = aE[X] \quad (2c)$
		Variance
		$\begin{aligned} \text{Var}[X] &= E[(X - E[X])^2] \\ &= E[X^2] - E^2[X] \end{aligned} \quad (3a)$
		$\text{Var}[aX + b] = a^2 \text{Var}[X] \quad (3b)$

## 2 Two Random Variables

### 2.1 Joint Distributions

Definitions:

$$F_{XY}(x, y) = p(X \leq x, Y \leq y) \quad (4a)$$

$$f_{XY}(x, y) = \frac{\partial^2 F_{XY}(x, y)}{\partial x \partial y} \quad (4b)$$

$$F_{XY}(x, y) = \int_{-\infty}^x \int_{-\infty}^y f_{XY}(s, p) dp ds \quad (4c)$$

Conditional distribution:

$$f_{Y|X}(y|x) = \frac{f_{XY}(x, y)}{f_X(x)}, \quad f_X(x) > 0 \quad (5a)$$

$$p(Y = y_j | X = x_k) = \frac{p(Y = y_j, X = x_k)}{p(X = x_k)}, \quad p(X = x_k) > 0 \quad (5b)$$

Expectation:

$$E[XY] = \int xy f_{XY}(x, y) dx dy \quad (6a)$$

$$E[g(X, Y)] = \int g(x, y) f_{XY}(x, y) dx dy \quad (6b)$$

$$E[X + Y] = E[X] + E[Y] \quad (6c)$$

Conditional expectation & Variance:

$$E[Y|X] = \begin{cases} \int y f_{Y|X}(y|x) dx dy \\ \sum_k y_k p(Y = y_k | X = x_j) \end{cases} \quad (7a)$$

$$E[X] = E[E[X|Y]] \quad (7b)$$

$$\text{Var}[Y|X] = E[Y^2|X] - E^2[Y|X] \quad (7c)$$

$$\text{Var}[Y] = \text{Var}[E[Y|X]] + E[\text{Var}[Y|X]] \quad (7d)$$

Independent random variables:

$$f_{XY}(x, y) = f_X(x)f_Y(y) \quad (8a)$$

$$F_{XY}(x, y) = F_X(x)F_Y(y) \quad (8b)$$

$$E[XY] = E[X]E[Y] \quad (8c)$$

$$E[g_1(X)g_2(Y)] = E[g_1(X)]E[g_2(Y)] \quad (8d)$$

$$\text{Var}[X + Y] = \text{Var}[X] + \text{Var}[Y] \quad (8e)$$

Marginal distribution:

$$f_X(x) = \int_{-\infty}^{\infty} f_{XY}(x, y) dy \quad (9a)$$

$$f_Y(y) = \int_{-\infty}^{\infty} f_{XY}(x, y) dx \quad (9b)$$

$$F_X(x) = F_{XY}(x, \infty) \quad (9c)$$

$$F_Y(y) = F_{XY}(\infty, y) \quad (9d)$$

### 2.2 Correlation, Covariance & Correlation Coefficient

- For two jointly-distributed random variables  $X$  and  $Y$ , covariance is given by

$$\begin{aligned} \text{Cov}[X, Y] &= E[(X - E[X])(Y - E[Y])] \\ &= E[XY] - E[X]E[Y]. \end{aligned} \quad (10)$$

Main covariance properties are:

$$\text{Cov}[X, X] = \text{Var}[X] \quad (11a)$$

$$\text{Cov}[X, Y] = \text{Cov}[Y, X] \quad (11b)$$

$$\text{Cov}[X, a] = 0 \quad (11c)$$

$$\text{Cov}[aX, bY] = ab \text{Cov}[X, Y] \quad (11d)$$

$$\text{Cov}[X + a, Y + b] = \text{Cov}[X, Y] \quad (11e)$$

$$\text{Cov}[X + Y, Z] = \text{Cov}[X, Z] + \text{Cov}[Y, Z] \quad (11f)$$

$$\text{Var}[X + Y] = \text{Var}[X] + \text{Var}[Y] + 2 \text{Cov}[X, Y] \quad (11g)$$

- Correlation coefficient (also termed as Pearson product-moment correlation coefficient) is given by

$$\rho_{XY} = \frac{\text{Cov}[X, Y]}{\sqrt{\text{Var}[X] \text{Var}[Y]}} \quad (12)$$

such that  $|\rho_{XY}| \leq 1$ .

- For two random vectors  $\mathbf{X} \in \mathbb{R}^m$  and  $\mathbf{Y} \in \mathbb{R}^n$ , the resulting  $m \times n$  covariance matrix is given by

$$\begin{aligned} \text{Cov}[\mathbf{X}, \mathbf{Y}] &= C_{\mathbf{XY}} \\ &= E[(\mathbf{X} - E[\mathbf{X}])(\mathbf{Y} - E[\mathbf{Y}])^T] \\ &= E[\mathbf{XY}^T] - E[\mathbf{X}]E[\mathbf{Y}]^T \end{aligned} \quad (13)$$

### 2.3 Relations

- When  $X$  and  $Y$  are *orthogonal*,  $E[XY] = 0$ .
- When  $X$  and  $Y$  are *uncorrelated*,  $\text{Cov}[X, Y] = \rho_{XY} = 0$ .
- When  $X$  and  $Y$  are *independent*, they are also uncorrelated (see also Eqs. 8).
- When  $X$  and  $Y$  are *jointly Gaussian* and uncorrelated  $\implies X$  and  $Y$  are independent.

## 2.4 Bi-variate & Multivariate Normal Distribution

Joint Gaussian distribution of  $X_1$  and  $X_2$

$$f_{X_1 X_2}(x_1, x_2) = \frac{1}{2\pi\sigma_1\sigma_1\sqrt{1-\rho^2}} \exp\left(-\frac{1}{2(1-\rho^2)} \times \left[ \frac{(x_1 - \mu_1)^2}{\sigma_1^2} + \frac{(x_2 - \mu_2)^2}{\sigma_2^2} - \frac{2\rho(x_1 - \mu_1)(x_2 - \mu_2)}{\sigma_1\sigma_2} \right] \right) \quad (14)$$

Multivariate Gaussian distribution of  $\mathbf{X} = (X_1, X_2, \dots, X_n)$  is given by

$$f_{\mathbf{X}}(\mathbf{x}) = \frac{1}{2\pi \det[\mathbf{C}_{\mathbf{X}}]} \exp\left\{-\frac{1}{2} (\mathbf{x} - \boldsymbol{\mu})^T \mathbf{C}_{\mathbf{X}}^{-1} (\mathbf{x} - \boldsymbol{\mu})\right\}, \quad (15)$$


---

## 3 Signal Characterization

### 3.1 Auto-signal

- Average:

$$E[x(t)] = \int_{-\infty}^{\infty} x f_x(x; t) dx \quad (16)$$

- Variance:

$$\text{Var}[x(t)] = E[x^2(t)] - E^2[x(t)] \quad (17)$$

- Auto-correlation

$$R_x(t_1, t_2) = E[x(t_1)x(t_2)] \quad (18a)$$

$$R_x(t_1, t_2) = R_x(t_2, t_1) \quad (18b)$$

$$R_x(t, t) = E[x^2(t)] \quad (18c)$$

- Auto-covariance

$$C_x(t_1, t_2) = E[x(t_1)x(t_2)] - E[x(t_1)]E[x(t_2)] \quad (19a)$$

$$C_x(t, t) = \text{Var}[x(t)] \quad (19b)$$

- Correlation Coefficient

$$\rho_x(t_1, t_2) = \frac{C_x(t_1, t_2)}{\sqrt{C_x(t_1, t_1)C_x(t_2, t_2)}} \quad (20a)$$

$$|\rho_x(t_1, t_2)| \leq 1 \quad (20b)$$

- When  $x(t_1)$  and  $x(t_2)$  are *orthogonal*,  $R_x(t_1, t_2) = 0$ .
- When  $x(t_1)$  and  $x(t_2)$  are *uncorrelated*,  $C_x(t_1, t_2) = \rho_x(t_1, t_2) = 0$ .
- When  $x(t_1)$  and  $x(t_2)$  are *independent*,  $R_x(t_1, t_2) = E[x(t_1)]E[x(t_2)]$ .

### 3.2 Cross-Signal

- Cross-correlation

$$R_{xy}(t_1, t_2) = E[x(t_1)y(t_2)] \quad (21)$$

- Cross-covariance

$$C_{xy}(t_1, t_2) = R_{xy}(t_1, t_2) - E[x(t_1)]E[y(t_2)] \quad (22)$$

- Correlation Coefficient

$$\rho_{xy}(t_1, t_2) = \frac{C_{xy}(t_1, t_2)}{\sqrt{C_{xy}(t_1, t_1)C_{xy}(t_2, t_2)}} \quad (23)$$


---

## 4 Wide-Sense Stationary (WSS) Process

Definition:

$$E[x(t)] = E[x(0)] = m_x = \text{const} \quad (24a)$$

$$R_x(t_1, t_2) = R_x(0, \tau) = R_x(\tau), \quad \tau = t_2 - t_1, \quad \forall t_1, t_2 \quad (24b)$$

## 4.1 Auto-signal

- Auto-correlation

$$R_x(\tau) = E[x(t)x^*(t + \tau)] = E[x(t - \tau)x^*(t)] \quad (25)$$

Properties:

$$R_x(-\tau) = R_x(\tau) \quad (26a)$$

$$R_x(0) = E[|x(0)|^2] = E[|x(t)|^2] \quad (26b)$$

$$\text{Var}[x(t)] = C_x(0) = \sigma_x^2 \quad (26c)$$

$$|R_x(0)| \geq R_x(\tau) \quad (26d)$$

Deterministic definition

$$R_x(\tau) = x(\tau) * x^*(-\tau) \quad (27)$$

- Auto-covariance

$$C_x(\tau) = R_x(\tau) - m_x^2 \quad (28)$$

- Correlation Coefficient

$$\rho_x(\tau) = \frac{C_x(\tau)}{C_x(0)} \quad (29)$$

- Power spectral density (PSD)

$$S_x(f) = \mathcal{F}\{R_x(\tau)\} \quad (30)$$

Properties:

$$S_x(f) = S_x(-f) \quad (31a)$$

$$S_x(f) \geq 0, \forall f \quad (31b)$$

$$S_x(f) \in \mathbb{R} \quad (\text{real numbers}) \quad (31c)$$

Average power

$$P_x = E[x^2(t)] = R_x(0) = \int_{-\infty}^{\infty} S_x(f) df \quad (32)$$

Deterministic  $X(f) = \mathcal{F}\{x(t)\}, X^*(f) = \mathcal{F}\{x^*(-\tau)\}$  definition

$$S_x(f) = X(f)X^*(f) = |X(f)|^2 \quad (33)$$

## 4.2 Cross-signal

- Cross-correlation

$$R_{xy}(\tau) = E[x(t)y^*(t + \tau)] \quad (34)$$

Properties

$$R_{xy}(\tau) = R_{yx}^*(-\tau) \quad (35a)$$

$$|R_{xy}(\tau)| \leq \sqrt{R_x(0)R_y(0)} \quad (35b)$$

$$|R_{xy}(\tau)| \leq \frac{1}{2} [R_x(0) + R_y(0)] \quad (35c)$$

Deterministic definition

$$R_{xy}(\tau) = x(\tau) * y^*(-\tau) \quad (36)$$

- Cross-covariance

$$C_{xy}(\tau) = R_{xy}(\tau) - m_x m_y \quad (37)$$

- Cross-PSD

$$S_{xy}(f) = \mathcal{F}\{R_{xy}(\tau)\} \quad (38)$$

Properties

$$S_{xy}(f) = S_{yx}(-f) = S_{xy}^*(-f) \quad (39)$$

Deterministic definition

$$S_{xy}(f) = X(f)Y^*(f) \quad (40)$$

## 4.3 White Noise Process

White noise process,  $n(t)$ , is WSS process that is characterized by

$$R_n(\tau) = \sigma^2 \delta(\tau) \quad (41a)$$

$$S_n(f) = \sigma^2 \quad (41b)$$

## 5 LTI and WSS Random Process

For LTI system with impulse response  $h(t)$

$$y(t) = x(t) * h(t) \quad (42)$$

Average

$$m_y = m_x \int_{-\infty}^{\infty} h(s) ds = m_x H(f=0) \quad (43)$$

### 5.1 Cross-correlation & cross-covariance

$$R_{xy}(\tau) = R_x(\tau) * h(\tau) \quad (44a)$$

$$C_{xy}(\tau) = C_x(\tau) * h(\tau) \quad (44b)$$

$$R_{yx}(\tau) = R_x(\tau) * h^*(-\tau) \quad (44c)$$

$$C_{yx}(\tau) = C_x(\tau) * h^*(-\tau) \quad (44d)$$

$$R_y(\tau) = R_x(\tau) * h(\tau) * h^*(-\tau) \quad (44e)$$

$$C_y(\tau) = C_x(\tau) * h(\tau) * h^*(-\tau) \quad (44f)$$

### 5.2 Power-Spectral Density (PSD) & Cross-PSD

Given frequency response  $H(f) = \mathcal{F}\{h(\tau)\}$ ,  $H^*(f) = \mathcal{F}\{h^*(-\tau)\}$

$$S_{xy}(f) = S_x(f) H(f) \quad (45a)$$

$$S_{yx}(f) = S_x(f) H^*(f) \quad (45b)$$

$$S_y(f) = S_x(f) H(f) H^*(f) = S_x(f) |H(f)|^2 \quad (45c)$$

Power of the process:

$$P_x = R_x(0) = \int_{-\infty}^{\infty} S_x(f) df \quad (46a)$$

$$P_y = R_y(0) = \int_{-\infty}^{\infty} S_x(f) |H(f)|^2 df \quad (46b)$$

## 6 Filtering of WSS Process

### 6.1 SNR

For an input

$$x(t) = s(t) + n(t), \quad (47)$$

where signal  $s(t)$  and noise  $n(t)$  are independent and  $E[n(t)] = 0$ , and output  $y(t)$

$$S_y(f) = S_x(f) |H(f)|^2 = S_s(f) |H(f)|^2 + S_n(f) |H(f)|^2, \quad (48)$$

where  $S_s(f) |H(f)|^2$  is signal output PSD and  $S_n(f) |H(f)|^2$  is noise PSD.

The input and output SNRs is given by

$$\text{SNR}_x = \frac{E[s^2(t)]}{E[n^2(t)]} = \frac{R_{ss}(0)}{R_{nn}(0)} = \frac{\int S_{ss}(f) df}{\int S_{nn}(f) df} \quad (49a)$$

$$\text{SNR}_y = \frac{\int S_{ss}(f) |H(f)|^2 df}{\int S_{nn}(f) |H(f)|^2 df}. \quad (49b)$$

### 6.2 Match Filter

The goal of filter  $h(t)$  is to provide maximum SNR at time  $t = t_0$  for *deterministic* signal  $x(t)$  and noise  $n(t)$ . The filter is given by

$$H(f) = \alpha \frac{X^*(f)}{S_n(f)} e^{-j2\pi f t_0}. \quad (50)$$

For white noise,  $n(t)$ , with  $S_N(f) = N_0/2$ , the filter is given by

$$H_{\text{mf}}(f) = X^*(f) e^{-j2\pi f t_0} \longleftrightarrow h_{\text{mf}}(t) = x(t_0 - t) \quad (51)$$

and the resulting maximum SNR is given by

$$SNR_{\max} = \frac{2}{N_0} \int_{-\infty}^{\infty} |X(f)|^2 df = \frac{2E_x}{N_0} \quad (52)$$

## 7 Poisson Process

The Poisson process,  $N(t)$ , is described by

$$p(N(t) = k) = e^{-\lambda t} \frac{(\lambda t)^k}{k!}, \quad k = 0, 1, \dots \quad (53a)$$

$$p(N(0) = 0) = 0 \quad (53b)$$

Time increment property

$$p(N(t_2) - N(t_1) = k) = p(N(t_2 - t_1) = k) \quad (54)$$

### 7.1 Campbell Theorem

Given the relation

$$y(t) = \sum_{k=1}^{\infty} g(t - t_k) \quad (55)$$

where  $t_k$  are Poisson event times and  $g(t)$  is causal system impulse response, resulting statistics is given by

$$E[y(t)] = \lambda \int_0^t g(u) du \quad (56a)$$

$$\text{Var}[y(t)] = \lambda \int_0^t g^2(u) du \quad (56b)$$

### 7.2 Auto-correlation & auto-covariance

$$C_y(t, t + \tau) = \lambda \int_0^t h(u)h(u + \tau) du \quad (57a)$$

$$R_y(t, t + \tau) = \lambda \int_0^t h(u)h(u + \tau) du + E[y(t)]E[y(t + \tau)] \quad (57b)$$

For time-limited  $h(t)$ , such that  $h(t) = 0, t > t_h$ , or at limit  $t \rightarrow \infty$

$$C_y(\tau) = C_y(t, t + \tau) \quad (58a)$$

$$R_y(\tau) = R_y(t, t + \tau) = \lambda \int_0^t h(u)h(u + \tau) du + E^2[y(t)] \quad (58b)$$

## 8 Different Supplementary Formulas

### 8.1 Derivatives

$$\frac{d}{dx}x^n = nx^{n-1}$$

$$\frac{d}{dx} \exp [f(x)] = \exp [f(x)] \frac{d}{dx}f(x)$$

### 8.2 Integrals

#### 8.2.1 Indefinite

$$\int x^n dx = \frac{1}{n+1}x^{n+1}, \quad n \neq -1$$

$$\int \exp(ax)dx = \frac{1}{a}\exp(ax)$$

$$\int x \exp(ax)dx = \exp(ax) \left[ \frac{x}{a} - \frac{1}{a^2} \right]$$

$$\int x^2 \exp(ax)dx = \exp(ax) \left[ \frac{x^2}{a} - \frac{2x}{a^2} + \frac{2}{a^3} \right]$$

$$\int \frac{1}{a^2+x^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a}$$

#### 8.2.2 Definite

$$\int_0^\infty \exp(-a^2x^2)dx = \frac{\sqrt{\pi}}{2a}$$

$$\int_0^\infty x^2 \exp(-a^2x^2)dx = \frac{\sqrt{\pi}}{4a^3}$$

$$\int_{-\infty}^\infty \delta(x)dx = 1$$

$$\int_{-\infty}^\infty f(x)\delta(x-a)dx = f(a)$$

### 8.3 Fourier Transform

#### 8.3.1 Properties

$$\frac{d^n}{dt^n} f(t) \xleftrightarrow{\mathcal{F}} (j2\pi f)^n F(f)$$

#### 8.3.2 Transform

$$u(t) \xleftrightarrow{\mathcal{F}} \frac{1}{2} \left( \frac{1}{j\pi f} + \delta(f) \right)$$

$$\exp(-at)u(t) \xleftrightarrow{\mathcal{F}} \frac{1}{a+j2\pi f}$$

$$t \exp(-at)u(t) \xleftrightarrow{\mathcal{F}} \frac{1}{(a+j2\pi f)^2}$$

$$\exp(-a|t|) \xleftrightarrow{\mathcal{F}} \frac{2a}{a^2+4\pi^2f^2}$$

$$\exp(-at^2) \xleftrightarrow{\mathcal{F}} \sqrt{\frac{\pi}{a}} \exp\left(-\frac{(\pi f)^2}{a}\right)$$

$$\cos(2\pi f_a t) \xleftrightarrow{\mathcal{F}} \frac{1}{2} [\delta(f-f_a) + \delta(f+f_a)]$$

$$\sin(2\pi f_a t) \xleftrightarrow{\mathcal{F}} \frac{1}{2j} [\delta(f-f_a) - \delta(f+f_a)]$$

### 8.4 Trigonometry

$$\sin^2(\alpha) = \frac{1}{2} (1 - \cos(2\alpha))$$

$$\cos^2(\alpha) = \frac{1}{2} (1 + \cos(2\alpha))$$

$$\cos(\alpha) \cos(\beta) = \frac{1}{2} [\cos(\alpha + \beta) + \cos(\alpha - \beta)]$$

$$\sin(\alpha) \sin(\beta) = \frac{1}{2} [\cos(\alpha - \beta) - \cos(\alpha + \beta)]$$

$$\sin(\alpha) \cos(\beta) = \frac{1}{2} [\sin(\alpha - \beta) + \sin((\alpha + \beta))]$$